

TAAD-5™ Triage Assessment For Addictive Disorders-5

The TAAD-5 is a brief structured interview designed to identify current alcohol and drug problems. This 10-minute assessment tool covers all DSM-5 criteria for substance use disorders. It offers high internal reliability. The TAAD-5 produces alcohol and drug use disorders profiles similar to those of more time intensive instruments.

A technician or paraprofessional can administer the TAAD-5 for interpretation by a qualified professional. A scoring summary is incorporated into each instrument and can be completed in 2-3 minutes.

Package of 25 • \$52.50

Item #: TA-T

Corresponding Manual


\$15.00 each • Item #: TA-M

 **The Change Companies®**

888-889-8866

www.changecompanies.net/products/popup.php?id=TA-T

Sample page



TAAD-5™
Triage Assessment for Addictive Disorders - 5
Norman G. Hoffmann, Ph.D.

Name: _____ Date: _____ Interviewer: _____
Age: _____ (1) Male (2) Female
Ethnic Background (check one): (1) Asian (2) African-American (3) Hispanic / Latino
(4) Native American (5) White / Caucasian (6) Biracial / Other
Highest Grade Completed (circle): 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16+

Darken the circle indicating the respondent's best answer to each question. Each question should be asked with regard to the past 12 months.

- In general, how often do you drink?
 - Daily
 - 4 to 6 days per week
 - 2 to 3 days per week
 - About once a week
 - At least 12 times a year
 - Less often
 - Never (*Go to # 3*)
- During the past 12 months, did you ever drink at least a fifth of liquor in one day? (That would be about 20 mixed drinks, three bottles of wine, or three six-packs of beer.)
 - No
 - Yes
 - 7 or more
 - 5 or 6
 - 3 or 4
 - 1 or 2
- Can you drink more now without feeling the effects than you once did?
 - No
 - Yes
- How often do you use other drugs?
 - Daily
 - 4 to 6 days per week
 - 2 to 3 days per week
 - About once a week
 - At least 12 times a year
 - Less often
 - Never (*Go to # 5*)
- Do you need larger amounts of drugs to get high than you once did?
 - No
 - Yes
- During the past 12 months, have you frequently used alcohol/drugs to relieve emotional discomfort, such as sadness, anger, or boredom?
 - No
 - Yes (alcohol only)
 - Yes (drugs only)
 - Yes (both alcohol and drugs)
- Have you frequently found yourself thinking about drinking/getting high?
 - No
 - Yes (alcohol only)
 - Yes (drugs only)
 - Yes (both alcohol and drugs)
- Have you occasionally had more to drink than you intended?
 - No (*Go to # 8*)
 - Yes
- How often would you say this happens?
 - Once a day
 - Several times a week
 - Several times a month
 - Several times a year
- During the past 12 months, have you set rules to limit your drinking or drug use that you failed to follow?
 - No
 - Yes (alcohol only)
 - Yes (drugs only)
 - Yes (both alcohol and drugs)

Copyright, 1995, 2012 Norman G. Hoffmann, Ph.D. All Rights Reserved. Distributed by The Change Companies, 888-889-8866, 755-886-2610. Reproduction or adaptation in any manner, in whole or in part, by any means, is a violation of copyright law and constitutes unethical and unprofessional conduct.

Applications:

- ◆ Any situation requiring fast, initial discrimination of likely a substance use disorder or no diagnosis.
- ◆ Ideal for preliminary triage to determine if a treatment referral or intake assessment is indicated.
- ◆ Other settings for use include:
 - a quick initial assessment of DUI/DWI offenders.
 - initial EAP assessments, mental health clinics, primary care settings or emergency rooms.

- initial field assessments by child protection, welfare workers or other social service providers.

Benefits:

- ◆ Offers a quick assessment of current substance use disorder criteria.
- ◆ Provides support for substance use disorder diagnoses in minutes.
- ◆ Documents negative findings for those individuals who deny problems.